Sl. No.:

	-	•	LO
Register Number			

2019 PHARMACEUTICAL CHEMISTRY (PG Degree Std.)

Time Allowed: 3 Hours

[Maximum Marks: 300

D.IPC/19

Read the following instructions carefully before you begin to answer the questions.

IMPORTANT INSTRUCTIONS

- 1. The applicant will be supplied with Question Booklet 15 minutes before commencement of the examination.
- 2. This Question Booklet contains 200 questions. Prior to attempting to answer, the candidates are requested to check whether all the questions are there in series and ensure there are no blank pages in the question booklet. In case any defect in the Question Paper is noticed, it shall be reported to the Invigilator within first 10 minutes and get it replaced with a complete Question Booklet. If any defect is noticed in the Question Booklet after the commencement of examination, it will not be replaced.
- 3. Answer all questions. All questions carry equal marks.
- 4. You must write your Register Number in the space provided on the top right side of this page. Do not write anything else on the Question Booklet.
- 5. An answer sheet will be supplied to you, separately by the Room Invigilator to mark the answers.
- 6. You will also encode your Question Booklet Number with Blue or Black ink Ball point pen in the space provided on the side 2 of the Answer Sheet. If you do not encode properly or fail to encode the above information, action will be taken as per Commission's notification.
- 7. Each question comprises four responses (A), (B), (C) and (D). You are to select ONLY ONE correct response and mark in your Answer Sheet. In case you feel that there are more than one correct response, mark the response which you consider the best. In any case, choose ONLY ONE response for each question. Your total marks will depend on the number of correct responses marked by you in the Answer Sheet.
- 8. In the Answer Sheet there are four circles (A), (B), (C) and (D) against each question. To answer the questions you are to mark with Blue or Black ink Ball point pen ONLY ONE circle of your choice for each question. Select one response for each question in the Question Booklet and mark in the Answer Sheet. If you mark more than one answer for one question, the answer will be treated as wrong. e.g. If for any item, (B) is the correct answer, you have to mark as follows:

- 9. You should not remove or tear off any sheet from this Question Booklet. You are not allowed to take this Question Booklet and the Answer Sheet out of the Examination Hall during the time of examination. After the examination is concluded, you must hand over your Answer Sheet to the Invigilator. You are allowed to take the Question Booklet with you only after the Examination is over.
- 10. Do not make any marking in the question booklet except in the sheet before the last page of the question booklet, which can be used for rough work. This should be strictly adhered.
- 11. Applicants have to write and shade the total number of answer fields left blank on the boxes provided at side 2 of OMR Answer Sheet. An extra time of 5 minutes will be given to specify the number of answer fields left blank.
- 12. Failure to comply with any of the above instructions will render you liable to such action or penalty as the Commission may decide at their discretion.

SEAL

1.	Cala	mine contains — and zinc	oxide	
	(A)	Ferric oxide	(B)	Ferrous sulphate
	(C)	Ferrous fumarate	(D)	Titanium dioxide
2.		limit test for arsenic is standardized ciple of this test is a modification of the		using special type of apparatus. The basic
	(A)	Sulphur test	(3)	Gutzeit test
	(C)	Lead test	(D)	Heavy metals test
3.	Unit	s of radioactivity is		
	(A)	Curie	(B)	Kg
	(C)	Jules	(D)	Mg
4.	Volu	ametric analysis is also called as		
	1	Titrimetric analysis	(B)	Qualitative analysis
	(C)	Gravimetric analysis	(D)	Semi quantitative analysis
5.		ntitative or semi quantitative test durity which is likely to be present is d		l to identify and control small quantity of as
	(A)	Quality control test		
	(7)	Limit test		
	(C)	Quantitative test		
	(D)	Identification test		
6.	Zinc	sulphate is assayed by		
	(A)	Non-aqueous titration		
	(B)	Gravimetry method		
	(C)	Precipitation titration		
	(0)	Complexometric titration	·	

7.	Whic	h of the following is used to relieve dental hypersensitivity?
grave .	(A)	Calcium phosphate
	(B)	Stannous fluoride
	10	Strontium chloride
	(D)	Sodium fluoride
8.	Dibas	sic calcium phosphate is used ———
	(A)	to treat dental caries
	(B)	to relieve dental hypersensitivity
	V.	as dentifrice
	(D)	to fill cavities
9.	The r	number of ligand donor atoms to which the metal is directly bonded is defined as
	(A)	Co ordination sphere
	(B)	Co ordination polyhedron
	VO	Co ordination number
	(D)	Counter ion
10.	Ethyl	lene diamine tetra acetic acid ionises in ———— stages
	(A)	1
	(B)	2
	(C)	3
* *	(1)	$oldsymbol{4}$
11.		——— is prepared by double decomposition reaction of hot ferrous sulphate and
	sodiu	m fumarate
	11)	Ferrous Fumarate
	(B)	Ferrous fluoride
	(C)	Ferrous glyconate
	(D)	Calcium Fumarate
DJP	C/10	
DOT.		그 나는 사람들은 사람들이 가장 하는 것이 되었다. 그 바람들은 그는 사람들은 사람들이 가장 하는 것이 되었다. 그런 그렇게 하는 것이 없는 것이 없는 것이 없는 것이 없는 것이 없는 것이 없다면 사람들이 없는 것이 없다면 없는데 없다면

14.	rerr	ic Ammonium citrate is assayed by —		
*	(A)	Iodometry titration	(B)	Precipitation titration
	(C)	Non-aqueous titration	(D)	Gravimetry method
13.	Whic	ch one of the following used as acid bas	e regu	lator
	(A)	Sodium carbonate	(B)	Sodium bi carbonate
	1	Sodium citrate tablets	(D)	Sodium hydroxide
		이 [캠프로그 실패] [인도 보고다] [인		
14.	Calc	ium disodium edetate is used in the —	27	
	1	treatment of lead poisoning	(B)	treatment of rheumatoid arthritis
	(C)	treatment of hyper acidity	(D)	treatment of constipation
15.	Sodi	um chloride infection is used as		
10.	(A)	Antacid	المحالة	Electrolyte replenisher
	(C)	Laxative	(D)	Antidote
	(0)			
16.	Codi	um thio sulphate is used as		
10.	(A)	Haematinic		Antidote for cyanide poisoning
	(A) (C)	Emetic	(D)	Expectorant
	(0)	Minetic	(D)	Expectorality
17.		nonium carbonate is used as	 .	
	(A)	Hematinic		
	(B)	Emetic Expectorant		
	(D)	Antidote	*	
	(D)	American		
10				na niman kalama
18.		ch one is the chelating agent among the	e optio	ns given below?
	(A)	Barium chloride		
	(B)	Silver Nitrate Ethylono diamino totro acetic acid	× 1	
	(D)	Ethylene diamine tetra acetic acid Potassium dichromate		
	(D)	rotassium dichromate		

19.	The	SI unit of co-efficient of viscosity is ————
	(A)	$ m kg~m^2S$
	(B)	$ m kg~m^{-1}S$
	(6)	$ m kg~m^{-1}S^{-1}$
	(D)	kg mS-
		등 이번 보는 방향 모양하다 하다 하네 사람들이 되는 사람들이 보는 사람들이 되었다. 그리는 사람들이 되었다.
20.	A liq	uid rises in a capillary tube. It is due to its ————
	(A)	Viscosity
	(B)	Vapour Pressure
	(C)	Density
		Surface tension
21.	Vice	osity of a liquid is a measure of ————
21.	(A)	Repulsive forces between the liquid molecules
	(A)	Frictional resistance
	(C)	Intermolecular force between the molecules
	(C)	그리다는 그 사람이 가다면 그녀를 가다고 하는 속하는 이 나는 사람들은 그리다는 생각하였다.
	(D)	Hydrogen bonding
		그 사람들이 내가 하다면 생각도 하는데 하다면 내가 되었다. 이 전에 되었다. 유럽 보다는 사람들
22.	Asso	ciation of molecule in water is due to ————
	(A)	Surface tension
	(B)	Viscosity
	Kor	Hydrogen bonding
	(D)	Optical activity
	1 100	마다는 네트리아 모양을 살이 되어 하는 그리다 경우에 되는 모든 다음 나는 보다 하다.
23.		greater the surface tension of the liquid, the higher is its capillary rise. This statement t suitable for ————
	(A)	Water
	9	Mercury
	(C)	Glycerin
	(D)	Acetic acid
		옷에 게 눈빛하게, 다느리가 되어 되어 뭐야? 이 아름이 느르지 않는 이번에는 목사였다.

24.	Perc	hloric acid is standadised using ——		
	(A)	Sodium carbonate		
	(B)	Potassium permanganate		
	(C)	Oxalic acid		
		Potassium hydrogen phthalate		
25.		——— is Aprotic solvent		
	(A)	Sulphuric acid		Chloroform
	(C)	Water	(D)	Acetic acid
26.	_	show deviations from Raoult's	Law	
,		Real solution	(B)	Ideal solution
	(C)	Super critical solution	(D)	Unsaturated solution
27.		states that, at constant ter	nperatur	e, the volume of a fixed mass of gas is
	inver	rsely proportional to its pressure.		
	(A)	Charles's law	(3)	Boyle's law
	(C)	Raoult's law	(D)	Beer's law
28.	Dum	a's method is used to determine —		
	(A)	halogens	(B)	sulphur
	(C)	phosphorous		Nitrogen
29.	Whic	ch one of the following methods is	used to	quantitatively determine the amount of
	ņitro	gen?		
	(A)	Rast method		
p =	(B)	Kjeldahl method		
	(C)	Zeisel's method		
	(D)	Herzig – Meyer method	Eig 16	

30.	Option called		e of polari	sation by equal and opposite amounts are
	(A)	Diastereo isomers		
	1	Enantiomers		
	(C)	Geometrical isomers		
	(D)	Cis-Trans isomers		
31.	or tw			or more different ways whether in one step nge is same no matter by which method the
	(1)	Hess law		
	(B)	Gibb's law		
	(C)	Law of mass action		
	(D)	Vant Hoff rules		
32.	Isome		l formula b	ut differ in arrangement of atoms in space
	4	Stereoisomers	(B)	Optical isomers
	(C)	Structural isomers	(D)	Chiral
33.	The h	neat of combustion can be determi	ned experi	mentally in a
	(A)	Polarimeter	(B)	Colorimeter
		Calorimeter	(D)	Refractrometer
34.	The called	그 그 그리는 살았는 것이 되고 그 그들은 경기에 가는 것 같아. 이 그리고 그는 것이 없는 것이 없는 것이 없는 것이다.	with the he	eat changes caused by chemical reaction is
	(A)	Phyto chemistry		
	The state of the s	Thermo chemistry		
	(C)	Photo chemistry		
	(D)	Electrochemical chemistry		
DJPC	2/19		8	

35.	The	The product of Molar Mass and specific refraction is called					
	(A)	Refractive Index	ж				
	(6)	Molar Refraction					
	(C)	Reflective Index					
	(D)	Molar Reflection					
36.	The	square of Refractive Index is used to	detect				
	(A)	Carbon bonds					
	(B)	Nitrogen bonds					
	(C)	Sulphur bonds					
	6	Hydrogen – bond complexes					
37.	Refr	active Index of water at room temper	ature is				
	(A)	1.55	(B)	1.44			
	1	1.33	(D)	1.22			
	· •						
38.	The	absorption co-efficients for dextro a	nd levo c	ircularly polarised light are different, this			
		rence is known as					
		Circular dichroism	(B)	Circular Polarity			
	(C)	Circular Asborptivity	(D)	Circular mobility			
39.	A m	athematical relation, connecting to	tal mola	r energy of fluid (gas or liquid) with its			
	volu	me and temperature is called as					
		Caloric Equations of state	(B)	Overlap repulsion force			
	(C)	Random packing modal	(D)	Cybotactic group model			
40.	The	average amount of energy required to	o dissocia	ate one mole is called as			
	(A)	endothermic compound		bond energy			
	(C)	exothermic compound	(D)	heat of reaction			
	s s "-						

41.	The	number of unpaired electrons in the ou	termo	st orbit is called as
	(A)	positive ions	(B)	negative ions
	(C)	valence electrons	(0)	free radicals
42.	Whi	ch one of the following reagent is used v	widely	in the preparation of alcohols?
	(A)	Volhard Reagent		
	(0)	Grignard Reagent		
	(C)	Benedict Reagent		
	(D)	Kolbes Reagent		
43.	Diaz	onium salts are used for preparations o	of ——	
	(1)	Dye stuffs	(B)	Aldehydes
	(C)	Ketones	(D)	Carboxylic acids
44.	m-Bi	romoToluene isomer is best synthesised	from	
	(A)	Bromination of toluene		
	(B)	Methylation of Bromobenzene		
	(6)	Diazotisation reaction with Toluene		
	(D)	From Nitro benzene		
45.	The	reaction of carboxylic esters with Grign	ard re	agent is an excellent method for preparing
	(A)	1° OH – (Primary alcohols)		
	(B)	2° OH – (Secondary alcohols)		
	VO	3° OH – (Tertiary alcohols)		
	(D)	Rectified spirit		
46.	Grigi	nard Reagent reacts with Water to give		
	W	Alkane	(B)	Alkene
	(C)	Alkyne	(D)	Acetylene
DJP	C/19	10		
	J. 10	10	87 B	그 사이 그 그 그 그 그 그 그 그 그 그 그 그 그 그 그 그 그 그

47.	Whic	Which one of the following has higher energies than the combining Atomic Orbitals (AOs)?					
	(A)	Bonding Molecular Orbitals (BMOs)					
	(6)	Anti Bonding Molecular Orbitals (AB)	MOs)				
	(C)	Atomic Molecular Orbitals (AMOs)					
	(D)	Linear Combination of Atomic Orbita	ls (LC	AOs)			
48.	Digox	kin is inhibitor of					
	(A)	Na+/K+ ATP-ase located in cardiac mu	ıltle				
	(B)	FADP Inhibtion in cell		하시스는 시간 회사가 가는 시간다			
	(C)	Ca ⁺ ATP-ase located in cardiac cells					
	(D)	Na ⁺ ATP-ase located in cardiac cells	Ū.				
		용하고 있는 항상이 하셨다고 있습니다.					
40	In 00	se of Digitalis purpurea, the cardiac ac	tivity i	is maximum with			
49.	(A)	Odoro side – H	(P)	Digoxin			
	(A)	Digitoxin	(D) .	Purpureo side – A			
•	(0)	Digitoxiii	(2)				
-0	XI/I :	h one of the following is 4–Quinolinone	. 3_Ca	rboxylic acid derivative?			
50.			(B)	Enoxacin			
	(A)	Nalidixic acid	(B)	Norfloxacin			
	(C)	Cinoxacin	(Normoxaciii			
51.	Piper	razine citrate is used in the treatment					
	(A)	Expectorant	(B)	Anti Tussive			
	10)	Anthelmintics	(D)	Anti Emetics			
52 .	Pyri	nidine nucleus is present in which of the	he follo	owing			
	(1)	Pyrantel	(B)	Niclosamide			
	(C)	Thio bendazole	(D)	Pyrazi Quantel			
53.	Diet	hyl carbamazine citrate comes under w	hich c	lass of Antholmintics			
	(A)	Benzimidazole	(B)	Nitro derivatives			
	(C)	Amides	(0)	Piperazines			

54. S-Enantiomer of ofloxacin is called as

(A) Spar floxacin

Levo floxacin

(C) Lome floxacin

(D) Balo floxacin

55. Which one of the following is an azole antifungal agent?

(A) Sordarin

(B) Butenafine

Ketoconazole

(D) Griseofulvin

56. The mechanism of action of Trimethoprim.

- (A) Blocks dihydrofolate reductase
- (B) Blocks t-RNA binding to m-RNA
- Blocks electron transport of bacteria
- (D) Blocks synthesis of dihydropteroic acid

57. The first marketed anti bacterial drugs were

(A) Cephalo sporins

(B) Pencillins

(C) Amoxy cillins

Sulfanilamide

58. Fill the Electrophilic aromatic substitution reaction:

$$(A) \xrightarrow{CH_3} \xrightarrow{H_2SO_4, SO_3} \xrightarrow{CH_3} + \xrightarrow{SO_3H} \xrightarrow{CH_3} \xrightarrow{CH_3} \xrightarrow{SO_3H} \xrightarrow{CH_3} \xrightarrow{CH_3} \xrightarrow{CH_3} \xrightarrow{CH_3} \xrightarrow{CH_3} \xrightarrow{CH_3} \xrightarrow{SO_3H} \xrightarrow{CH_3} \xrightarrow{$$

59. Which of the following antifungal should not be used during pregnancy?

- (A) Isavuconazonium sulfate
- (B) Nafti fine

(C) Butena fine

(D) Terbena fine

60.	Whic	th of the following is a pyrimidine of	derivative?	
	(A)	Proguanil HCl		
		Pyrimethamine		
	(C)	Cycloguanil Pamoate		
	(D)	Chlorophenyl guanidine		
61.	What	t could be the starting matial for th	ne synthesi	is of Phenobarbital?
		Benzyl chloride		
	(B)	Ethyl Methyl Ketone		
	(C)	Phenyl Acetamide		
	(D)	Phenyl Acetyl Chloride		
62.	Trim	ethoprim exhibits — wh	nich activit	y ?
	(A)	Antihistaminic		
	(B)	Anthelmintics		
		Anti bacterial		
	(D)	Anti depressant		
63.	Whic	h one of the following is osmotic di	uretic used	d in treatment of hypertension?
	(A)	Mannitol	(B)	Sorbitol
	(C)	Acetazolamide	(D)	Chlorthiazide
64.	Meta	zocine is unsuitable for usage beca	use of its	
		Psychomimetic side effects	(B)	Neurologic side effects
	(C)	Nephrologic side effects	(D)	Hepatic side effects

65.	Para	ıldehyde is used as		
/.	(A)	Anti depresents		
		Sedative		
	(C)	Anti convulsant		
	(D)	Anti histamine		
				보는 되면 하고 있는 아이트 사람들이 되었다.
66.	Thio	pental sodium is administered throu	ıgh ——	route.
	(A)	INNALATCON		
	(0)	INTRA VENOKS		
	(C)	ORAL		
	(D)	TOPICAL		
67.		eine is derived from morphine by dis	placemer	nt of the hydrogen atom of the phenolic-OH
	(A)	$-NH_2$		
	(B)	$-\mathrm{C}_2\mathrm{H}_5$		
	(C)	-СНО		
	10)	$-\mathrm{CH}_3$		
68.	Sulp	honamids are generally used to prod	luce ——	effect
	(1)	Antibacterial		
	(B)	Antibiotics		
	(C)	Anti malarial		
	(D)	Antihistamine	The Automotive	
69.	In th	ne Phenobarbital aray with N/10n N	aOH Phe	roharbital acts as
	(A)	Weak Mono basic acid	(B)	Weak bibasic acid
6 ·	(C)	Strong acid	(D)	ester
70.	Mafe	enide belongs to the category of which	h nucleus	s?
	(1)	Sulphonamide	(B)	STEROID
	(C)	PHRIDINE	(D)	PHENANTHARENE
DJP	C/19	그런 하는 나는 내가 가게 가는 것이다.	14	

71.	VVIIIC	in one of the following is a CNS still the	and an	aloid
	(A)	Guinine	(B)	CINCHONINE
	10	STRYCHNINE	(D)	Ephedrine
72.		refers to the specific type of	of instr	umentation where in the molar elipticity of
	an op	otically active substance is measured		
	(A)	Optical rotary dispersion	(B)	IR Spectrometer
		Circular dichroism	(D)	Mass spectrum
73.	TCA	cycle is commonly known as ———	-	
	(A)	Ko Warburg-Lipman Pathway	(B)	.Kreb's cycle
	(C)	EMP Pathway	(D)	Cori cycle
\				
74.	A sol	lution of cholesterol in chloroform w	hen tre	eated sulphuric acid and acetic anhydride
	gives	a green colour. This reaction is called	l ——	
	(A)	Salkowski reaction	(B)	Mayer's reaction
	B	Libermann-Burchard reaction	(D)	Hager's reaction
75 .	Oest	rone may be reduced to —————	— by ca	atalytic hydrogenetion, by LiAlH ₄
	(A)	Oestrogen	(B)	androgen
	(C)	Oestrotriol	(D)	Oestrodiol
76.	Cort	isone is used in the treatment of		
	(A)	Rheumatiod arthritis	(B)	Goitre
	(C)	Heart disease	(D)	Diabetis mellitus

77.	Quii	nine on controlled oxidation with chrom	ic aci	d gives
	M	Quininic acid and meroquinene		그는 일반 시간의 기일 있었다.
	(B)	Laiponic acid and quinine acids		
	(C)	Quinine acids and 4-methyl-6-methox	ky qui	noline
	(D)	Mosoquinene and laiponic acid		
78.	The	nature of carbon skeleton in a Nitrogen	eous	heterocyclic ring is determined by
	(A)	Zerehinoff's method		
	(B)	Zeisol's method		
	(C)	Clarke's method		
	VI)	Hofmann's exhaustive methylation m	ethod	
79.	The	number of asymmetric carbon counters	in Ep	hedrine is
	(1)	2	(B)	4
	(C)	8	(D)	16
80.	Trop	oine and pseudotropine are optically —		because of
	(A)	inactive, absence of chiral centres		
	(B)	active, two chiral centres		
	(C)	active, internal compensation		보고하다가 됐다면 맛이 없는 선생님
	100	inactive, internal compensation		
81.	Nico	tine when oxidised with dichromate-sulp	ohuri	c acid, if forms
	W	Nicotinic acid	(B)	Pyridine – 2 – carboxylic acid
	(C)	Pyridine – 4 – carboxylic acid	(D)	Benzoic acid
82.	The p	presence of N-methyl group and their nu	ımbei	rs may be determined by means of
	(A)	Hofmann's exhaustive methylation me		
	(B)	Van Braun's method		
	1	Herzig-meyer method	. *	
	(D)	Emde degradation method		
		그는 그는 그는 이 이 이 이 그는 것이 되었다. 그는 그 이 이 이 이 이 이 이 이 이 이 이 없어 그는 것이다. 그는 것이 없는 것이다. 그는 것이 없어 그렇게 되었다. 그는 것이다. 그는		

83.	The	blue shift means		너 뭐하다. 이 시네티얼마 나라하다
		A shift of λ_{max} to shorter wavelength		
	(B)	Increase in the intensity of an absorpt	ion	
	(C)	A shift of λ_{\max} to longer wavelength		
	(D)	Decrease in the intensity of an absorpt	tion	
84.	In o	prism monochromator the working princ	pinlo	io
04.	ma	prism monochromator the working princ	ipie	15
	(A)	Reflection	(B)	Scattering
	(C)	Re-inforcement		Dispersion
85.	Abso	rption of light in the ultraviolet regions	of the	e spectrum is due to the presence of a/an
	(A)	σ -electrons		chromophore
	(C)	auxochromes	(D)	electrolytes
· · · ·				
86.	-	law is defined as the intensity	y of a	a beam monochromatic radiation decreases
	expo	nentially with the number of absorbing r	mole	cules.
	A)	Beer's	(B)	Lambert's
	(C)	Brag's	(D)	Hess
87.	Quin	nine is highly fluorescent in 0.05 m sul	phur	ic acid but not in 0.1 m hydrochloric acid
	beca	use of		
	(1)	Collisional quenching		
	(B)	Static quenching		
	(C)	Tyndall scatter		
	(D)	Rayleigh scatter	r	

88.	The	efficiency of a chromatography columi	ı is mea	asured by its number of
	(A)	Elution	(8)	Theoretical plates
	(C)	Mobile phase	(D)	Compounds in mixture
89.	The	upper surface of a column should be p	rotecte	d by using ———— in gel filteration.
	(A)	Adsorbents	(B)	Charcoal
		Filter paper	(D)	Absorbents
90.	Mech	nanism of separation in TLC of paraffi	n oil or	silicon oil coated on silica is
i e	(1)	Reversed phase partition	(B)	Adsorbtion
	(C)	Reversed phase absorbance	(D)	Ion exchange
			, X	
91.	*Whic	h of the following developing reagen	it is us	ed for visualization of amino acid in thir
	layer	chromatography?		
	(A)	Iodine vapour	(B)	Bratton-Marshall reagent
		Ninhydrin reagent	(D)	Dragendroff's reagent
92.		can be made visible through	exposur	re of the TLC plate to iodine vapor.
	(A)	electrolytes	(B)	organometallic compounds
	VI	organic analytes	(D)	inorganic analytes
93.	The r	nost common iodine isotope used for b	iologica	al arrays
	(A)	131 _I	(B)	$^{125}\mathrm{I}$
	(C)	¹²⁴ I	(D)	136 _I

94.	IR al	osorption spectra are due to changes	in ——	energy accompanied by change in
	rotat	ional energy.		
	(A)	electronic		
		vibrational		
	(C)	nuclear spin		
	(D)	molecular charge		
95.	In IF	R, two atoms (non bonded) connected	d to a cer	ntral atom move up and move down below
	the p	lane is called ———— vibration		
	(A)	Twisting	(B)	Scissoring
	W/S	Wagging	(D)	Rocking
96.	A cor	mmon detector employed to detect IF	R radiatio	on is the
	(A)	Photovoltanic cell	(B)	Photomultiplier
	(C)	Crystal	VA .	Thermocouple
97.	Defo	rmation vibrations in IR spectroscop	y is calle	${ m ed}$ as
		bending vibrations		
	(B)	symmetric vibrations		
	(C)	asymmetric vibrations		
	(D)	stretching vibrations		
98.	In N	MR, the inter action between differen	ent hydro	gens in a molecule is
	(A)	chemical shift	(B)	coupling constant
	(C)	spin-spin coupling	(D)	deshielding

33.	Sour	um carbonate added to not solution o	1 pnosp	noric acid gives
	(1)	Disodium hydrogen phosphate	(B)	Sodium phosphoric acid
	(C)	Sodium bicarbonate	(D)	Sodium acid phosphate
100.	-	is dimethyl polysiloxane of g	rade 20	0
	(A)	Sulphurated potash		Dimethicone
	(C)	Potash	(D)	Kaolin
101.	-	are used for removing toxic s	ubstan	ces from GIT, caused due to poisonina or in
	diarr		,	
	(A)	Acidifiers	(B)	Adsorbents
	(C)	Antacids	(D)	Laxatives
102.	From	the following identify the chemical fo	ormula	
	(A)	$\mathrm{mg_6Al_2(OH)_{16}CO_3.4H_2O}$	(3)	$\text{Al}_5 \text{mg}_{10} (\text{OH})_{31} (\text{SO}_4)_2$
	(C)	$Al_2 mg_6 (CO_3)_2 (OH)_{14}.4H_2O$	(D)	${ m CaCO_3}$
103.		——— is obtained when magnesium	chloric	le reacts with sodium hydroxide
	(A)	Magnesium phosphate	(B)	Manganese oxide
	(C)	Aluminium hydroxide	(6)	Magnesium hydroxide
104.	Epsoi	m salt is known as		
	(A)	Calcium carbonate	(B)	Calcium hydroxide
	5	Magnesium sulphate	(D)	Magnesium carbonate
105.	Roche	elle salt is known as		
	(A)	Sodium sulphate	(3)	Sodium potassium tartrate
	(C)	Sodium phosphate	(D)	Bismuth sub carbonate
	4.56			

106.		h of the following is used to preve- test for Iron?	nt the pre	cipitation of Iron as ferric hydroxide in th
	(A)	Fumaric acid	(B)	Acetic acid
		Citric acid	(D)	Tartaric acid
107.	Magr	nesium sulphate is uses as		
	(A)	Antacid	(B)	Saline purgative
	(C)	Electrolyte replemishes	(D)	Dental product
108.	Whic	h of the following injections is used	l for the di	agnosis of hematological disorders?
	(A)	Gold (198Au) injection		
	(B)	Cyanocobalamin (60Co)		
		Ferric citrate (59Fe) injection		
	(D)	Sodium iodide (131I) injection		
109.	Sodiu	um orthophosphate solution is used	l in the —	
	(A)	Study of sodium exchange		
	(B)	Extra cellular water measuremen	nt	
	(0)	Treatment of polycythemia		
	(D)	Determination of myocardial bloc	od flow	
110.	Rubio	dium chloride injection is used in t	he ———	
•	(A)	determination of myocardial bloo	d flow	
	(B)	study of thyroid uptake		
	(C)	treatment of polycythemia		
	(D)	study of potassium exchange		
111.		is a material used for clean	ing of teet	h and adjacent gums
	(A)	Dental caries		
	(B)	Oral Antiseptic		
	(0)	Dentifrice		
	(D)	Dental hypersensitivity		나는 하는 사람이 없는 그래?

112.		———— can be obtained by careful n	eutraliza	tion of hydrochloric acid with lime.
	(A)	Calcium gluconate		Calcium chloride
	(C)	Potassium chloride	(D)	Sodium chloride
113.	Assa	y of Ammonium chloride is by		
	(A)	Complexometry	(B)	Non-aqueous titration
	(C)	Acidimetry	(6)	Modified Volmard's method
114.	Calc	ium hydroxide is assayed by		
	(A)	Acidimetry	(B)	Alkalimetry
		Complexometric titration	(D)	Non-aqueous titration
115.		ne assay of aluminium hydroxide ge in which conditions only the complex	The state of the s	is added to maintain an alkalin on is complete.
	(A)	Disodium edetate		
	(B)	Ammonia ammonium chloride		
		Hexamine		
	(D)	Magnesium oxide		
116.	In co	ompound sodium chloride solution, so	dium is	determined by ————.
	(A)	Spectro photometry		
	(B)	Flame photometry		
	(C)	Fluorimetry		
	(D)	Turbidometry		
117.	Calci	ium gluconate is assayed by ———	 .	
	(A)	Precipitation titration		
	(B)	Non-aqueous titration		
	(C)	Gravimetry method		
	(1)	Complexometric titration		회사인 하게 가득하다는 것 같아.

118.	Whic	h of the following is used as oral antiseptic?
	(A)	Sodium perborate
	(B)	Sodium Thio Sulphate
	(C)	Sodium Sulphite
	(D).	Sodium Methoxide
119.		has been the traditional cleaning-polishing agent for most tooth pastes and
	tooth	powders
	(A)	Sodium Carbonate
	-OS)	Calcium Carbonate
	(C)	Calcium Sulphate
	(D)	Sodium bi carbonate
120.	Calci	um chloró hypochlorite is known as ———
	(A)	Epsom salt
	(B)	Precipitated chalk
	(0)	Bleaching powder
	(D)	Lime salt
1 m		
121.	Britis	sh anti lewisite (B.A.L) is
	(A)	Leucovarin calcium
	(B)	D. Penicillamine
		Dimencaprol
	(D)	Editic acid
122.	The S	SI unit of surface tension is ———
	(A)	dyne cm
	(B)	$ m dyne^{-1}~cm$
	(C)	Nm
	-5	Nm^{-1}

125.	Gly	cerol has an unusually high viscosity	mainly	because of its high capacity to form
	(A)	Free radicals	(B)	Ionic bonds
	WAY.	Hydrogen bonds	(D)	Binary compounds
124.	As t	he temperature of a liquid increases,	its visco	eity
	(A)\$		(B)	increases or decreases
	(C)	remains same		decreases
	(0)	Temams same	. (20)	uecreases
125.	Osn	notic pressure of a solution is a/an —		
	(A)	Colloidal property	(B)	electrochemical property
		colligative property	(D)	catalytic property
126.	In a	cetone – chloroform system the deviat	ion from	Raoult's law is ———
	(A)	positive	10)	negative
	(C)	zero	(D)	positive and negative
127.	In m	ass spectra the most intense peak is l	known a	s
	(1)	Base peak	(B)	Hydrocarbon peak
	(C)	Fragment ion peak	(D)	Rearrangement peak
128.	Peop (A)	ble stranded in lifeboats on the ocea Surface tension is too low	an canno	ot drink the seawater. The reason is its Osmotic pressure is too high
	(C)	Viscosity is too high	(D)	Freezing temperature is too high
129.	Whic	ch one is a colligative property		
•		Osmotic pressure	(B)	Molecular weight
	(C)	Surface tension	(D)	Atomic volume

130.	Nitro	ous oxide is manufactured by the action of heat on
	(A)	Sodium nitrate
	(B)	Potassium nitrate
		Ammonium nitrate
	(D)	Ammonium carbonate
,		
131.	In te	sting oxygen for carbon di-oxide, the gas is passed slowly through a 3% solution of
	(A)	Calcium chloride
	(B)	Magnesium chloride
		Barium hydroxide
	(D)	Ammonium hydroxide
132.	The	change in enthalpy that take place when one mole of the compound is farmed from its
	elem	ents. If is usually represented by
	(A)	ΔH_{f} (B) ΔH
	(C)	$\Delta^{\circ}\mathrm{H}^{\circ}$ (D) $\Sigma\mathring{\Delta}\mathrm{H}^{\circ}$
199	Tho	separation of racemic modification into enantiomers is called
133.		Revolution
	(A) (B)	Regression
		Resolution
	(D)	Recession
	(D)	recession
134.		——— equation gives variation of partial vapour pressure of the constituents of a liquid
134.	mixt	ure with the variation of the composition in the liquid phase
	(A)	Nernst (B) Gibb's
		Duhem-Margules (D) Raoult's
		그 경기는 그 생물이 그 사람이 가면 가장하는 사람들이 있는 것이 되었다. 그 사람들이 되었다면 가장 그는 그 가게 하는 것은 것이다.

135.	A car	bon atom which is bonded to four different group is called as
	(1)	Asymmetric carbon atom
	(B)	Symmetric carbon atom
	(C)	Optical isomerism
	(D)	Geoisomerism
136.		h equation is the basis for the relationship between voltage generated and relevant entration at each electrode?
	(A)	Polynomial equation
	(B)	Simultaneous equation
		Nernst equation
	(D)	Brag's equation
137.	A spe	cial type of functional isomerism in which the isomers are in dynamic equilibrium with other
	(A)	Metamerism
	B) .	Tautomerism
	(C)	Enantiomers
	(D)	Optical isomers
		그래, 그는 이 말이라는 먹다. 나는 말이들이 아니라지만, 그렇다 이상 그리라는 그리다.
138.	The p	phenomenon in which one of the products itself acts as a catalyst is known as
	(A)	Positive Catalysis
	(B)	Negative Catalysis
	S	$\operatorname{Auto}-\operatorname{Catalysis}$
	(D)	Promoter
		엄마하는 나는 바다 그 사람이 하네요? 그리다는 그리는 그리는 그리고 하는 것이 없다.
139.		angle of rotation of the plane polarized light produced by a liquid with a solution of ne 1 ml and 1 gm of substance with 1 dm length is called as
•	(A)	Optical activity
	(P)	Specific rotation
	(C)	Rotatory action
	(D)	Specific absorbance
DJPC	/19	26

140.		change in enthalpy that takes place we ent is defined as	hen on	e mole of the compound is formed from its
		Heat of formation	(B)	Heat of solution
	a(C)	Heat of combustion	(D)	Heat of neutralisation
141.	The	temperature at which the two conjug	rate sol	ution merge into one another to form one
141.		r is called as	,000 001	
	(A)	conjugate system	(B)	critical minimum temperature
	(C)	tie line	1	critical solution temperature
142.		n a — aromatic amine is di sodium nitrite, Diazonium salts are fo		in cold aqueous mineral acid and treated
	(A)	Primary	(B)	Secondary
	(C)	Tertiary	(D)	Quarternary
143.	Vitar calle		th antii	mony trichloride gives Blue colour. This is
	$\langle A \rangle$	Carr-Price test	(B)	Wagners test
	(C)	Fehling test	(D)	Borntragers test
144.	The 1	unit for dipole moment is		
	(A)	amu	(B)	esu
	16	debye, D	(D)	centipoise, CP
145.	LCA	O' stands for ———.		
110.	(A)	Least Combination of Atomic Orbita	ıls	
	(B)	Last Combination of Atomic Orbital		
		Linear Combination of Atomic Orbit	tals	
	(D)	Largest Combination of Atomic Orb	itals	
146.	The	unit 'amu' means		
140.	(A)	Average mass unit	(B)	Average molecule unit
		Atomic mass unit	(D)	Atomic molecule unit
100	(TATOMAN AND MAKE	(/	

147.	Hete	erocyclic compounds are mainly synthes	sised f	rom
	(A)	Dimethyl Sulfoxide (DMSO)		
	(B)	Tri Fluoro Acetic acid (TFA)		
	(C)	Poly Ethylene Glycol (PEG)		
	1	Ethyl Aceto Acetate (EAA)		
148.	Vita	min-K3 is called as		
	(A)	Farnoquinone	(B)	Menaquinone
	S	Menadione	(D)	Phylloquinone
				마루 : 1 시간 1 시간 : 1 - 1 시간 : 1 시
149.	Ethy	l aceto acetate can be synthesised by		
	(A)	Aldol condensation		
	16)	Claisen condensation		**
	(C)	Clemmenon reduction		
	(D)	Birch Reduction		
150.	Alky	lated derivatives of acetic acid are obtain	ined f	rom — by acid hydrolysis.
	(A)	Malonic Esters		
	(B)	Succinic Esters		
	(C)	Diethyl Esters		
	()	Aceto Acetic Esters		
151.	CH_3	$-\mathrm{CO}-\mathrm{CH}_2-\mathrm{COOC}_2\mathrm{H}_5$ is called as		
	(A)	Diethyl Malonate (DEM)	(B)	Ethyl Aceto Acetate (EAA)
ė.	(C)	Tri Ethyl Amine (TEA)	(D)	Diethyl Amine (DEA)
	· V			
152.	Mole	cules like $ mH_{2}, O_{2}, N_{2}, Cl_{2}$ and $ mBr_{2}$ have		——— Dipole moments.
	(A)	Very high		Very low
	1	Zero	(D)	High

153.	In a	E1 reaction involving an altyl halide and a base, the rate of the reaction is
	(A)	linearly depends on the concentration of the alkyl halide only
	(B)	linearly depends on the concentration of both reactants
	(C)	independent of the concentration of the alkyl halide
	(D)	is independent of the concentration of both reactants
154.	The	major product of E2 reaction of alkyl fluorides is the ———————————————————————————————————
	(A)	Terminal
	(B)	More stable
	6	Less stable
	(D)	Symmetric
155.	2-Br	omo butane heated with alcoholic KOH gives
	(A)	Cyclobutans
	(B)	1 Butanol
•	10)	1-Butene and 2-butene
	(D)	Tri substituted butune
156.	Whic	h reaction takes place with inversion of configuration?
	(A)	SN^1 reaction
	(B)	SN ² reaction
	(C)	Asymmetric synthesis
	(D)	Stereo selective reaction
157.	Elect	rophiles are
	(A)	Electron rich species
	(6)	Electron deficient species
	(C)	Neutrons rich species
	(D)	Proton deficient species

158.	Tri n	nethoprim and sulfonamide combina	ation give			
	$\langle A \rangle$	Synergistic action				
	(B)	Reversible Antogonist				
	(C)	Antagonistic action				
	(D)	Irreversible Antagonist				
159.	The 1	nost serious adverse effect associat	ed with p	yrazinamides is		
	(A)	Cyto toxicity				
	(B)	Hepato toxicity				
	(C)	Nephro toxicity				
1	(D)	Neuro toxicity				
160.	Whic	h of the following is Amides contain	ning Anth	elmintics?		
	(A)	Furoesimide				
	1	Niclosamide				
	(C)	Actimide				
	(D)	Benzamide				
161.	Albendazole contains which of the following nucleus?					
	(A)	Pyrazole				
	(B)	Benzimidazole				
	(C)	Indole				
	(D)	Quinoline				
162.	Resor	nance in Benzene is due to				
	1	Delocalisation of π –electrons	(B)	Stable π -electrons		
	(C)	Stable σ -electrons	(D)	Unstable σ -electrons		
163.	Whic	h one of the following is a Benzimid	lazole Ant	chelminthic?		
	(A)	Piperazine Citrate		Mebendazole		
	(C)	Prazi Quantel	(D)	Avermectin		

- 164. Barbiturate and Benzodiazepines (Sedative and Hypnotics) are
 - (A) GABA receptor Agonist
- (B) GABA receptor Antagonist
- (C) DUPA receptor Agonist
- (D) DUPA receptor Antagonist
- 165. Which one of the following does not have asymmetric carbon?
 - (A) Halothane

(B) Isoflurane

(C) Desflurane

- (d) Methoxyflurane
- 166. Which one of the following Antidepresent is selective Nor epinephrine Receptor Inhibitor?
 - (A) Citalopram

(B) Sertraline

Desipramine

- (D) Fluoxetine
- 167. Which form of Triprolidine is pharmacologicaly active?
 - (A) CIS-form

(3) Trans-form

(C) R-form

- (D) S-form
- 168. Choose the correct chemicals structure of Dapsone

$$(A) \quad \left\langle \bigcirc \right\rangle - \left\langle \begin{matrix} 0 \\ \parallel \\ \parallel \\ 0 \end{matrix} \right\rangle$$

(B) $CH_3 \longrightarrow CH_3$ O

CH₃
O

CH₃

- O_2N \longrightarrow S \longrightarrow NO_2
- 169. A cardioselective β –adrenergic blockers is
 - (A) Nitroglycerin

B) Propranolol

(C) Verapamil

(D) Bepridil

170.	The	The drug ketamine is used as ————						
		Anaesthetic	(B)	Anti-tubercular agents				
	(C)	Anti histamine	(D)	Anthelmirtics				
171.	Whi	ch one of the following is a Thiophene	deriva	tive				
		methapyrilene hydrochloride						
	(B)	meclizine hydrochloride						
	(C)	Buclizine hydrochloride						
	(D)	Chlorcyclizine hydrochloride						
172.	Which form of Atomoxetine is more active Anti depressant than other form?							
		R-Atomoxetine						
	(B)	S-Atomoxetine						
	(C)	RS-Atomoxetine						
	(D)	Cis-Atomoxetine						
173.	Mart	essentiaum consists of the grain of	the ce	real ——— belongs to the family				
	Gran	ninea						
	(A)	D-glucose	(3)	Barley				
	(C)	O-galactose	(D)	melibiose				
174.		is a pro-vitamin D ₂ which i	s found	d both in plants, animals and yeast.				
	(A)	Lansesteral	(B)	Stigmasterol				
		Ergosterol	(D)	Bile acid				
	- Carrier - Lan							

175.	w nic	on of the following is pyrrollaine alkalo	ια:		
	(A)	Ricinine			
	(B)	Coninine			
	S	Hygrine			
	(D)	Reserpine			
					5
176.	Molis	sch test is used for the identification of			
	(A)	Proteins			
	B)	Carbohydrates			
	(C)	Alkaloids			
	(D)	Steroids			
177.	Chole	esterol contains — nu	ımber	of carbon atoms.	
		27	(B)	17	
	(C)	24	(D)	28	
					*
178.	Whic	h of the following amino acids has a ph	nenolio	c ring?	
	(A)	Proline	(B)	Alanine	
	(C)	Prytophan	(Tyrosine	
179.	The r	number of chiral centres in Glucose is			
	(A)	8		4	
	- Da - 11 1				

100.	wan	tose on hydrolysis by dilute acids yields		
	(A)	two molecules of fructose		
		two molecules of D-glucose		
	(C)	one molecules of D-glucose and one m	olecu	le of fructose
	(D)	three molecules of D-glucose		
181.	Ligh	nt source used for the measurement in th	he ult	raviolet region is a
	(A)	Tungsten filament lamp		
	(B)	Denterium discharge lam		
	(C)	Globar rod		
	(D)	Nernst glowers		
182.	Whic	ch of the following reduces the fluoresce	nce of	riboflavine by static quenching?
	(A)	EDTA	(B)	Dimercaprol
	(C)	Penicillamine	(1)	Caffeine
183.	Grad	des of silica used in HPTLC has the part	ticle s	ize as ———— and ———.
	(A)	large and uniform		
	(B)	small and uniform		
	(C)	large and irregular		
	(D)	small and irregular		
184.	Whic	ch of the following is used for the determ	ninati	on of molecular weight?
	(A)	Gas chromatography	(B)	Paper chromatography
	1	Gel filtration	(D)	Ion-exchange

185.	Quar	ntum yield of fluorescence would be equa	ar to					
	(A)	number of photons emitted - number of	of pho	otons absorbed				
	(B)	photons absorbed – photons emitted						
		number of photons emitted / number of	f pho	tons absorbed				
	(D)	number of photons absorbed / number	of ph	otons emitted				
186.	The	quantum efficiency fluorescence decreas	es wi	th increasing				
	(A)	viscosity	0	temperature				
	(C)	pH	(D)	pressure				
107	HDI	C – state silica consists of porous microp	oortio	les with a ———— (or) ———				
187.	shap		Jartic	les with a				
	(A)	Spherical (or) Regular	(B)	Non spherical (or) Regular				
	1	Spherical (or) Irregular	(D)	Non spherical (or) Irregular				
188.	The	The mobile phase in Reverse-Phase HPLC comprises — and — and —.						
100.	(A)	water and petroleum ether	r					
		water and methanol						
	(C) ·	water and carbon tetrachloride						
	(D)	water and cyclohexane						
	(2)							
100	T TT	DIC the efetien are phase is polar and	tho.	mobile phase is non-polar, then it is called				
189.	In H		the	mobile phase is non-polar, then it is cance				
	(A)	Normal-phase partition						
	(B)	Reversed-phase partition		시시다리는 학생이 지하다는 생각이				
	(C)	Ion-pair						
	(D)	Ion-exchange						
190.		IPLC, while supplying mobile phase by ired to smoothout the pulses.	y me	chanical pump, a — device is				
	(A)	Gauze		Damping				
	(C)	Temperature	(D)	Injection				

191. In an applied magnetic field in NMR study, the number of orientations of a r				number of orientations of a nucleus with		
	spin	number I, is given by the formula of	of			
	(A)	2(I+1)	(B)	I+1 .		
•		2I+1	(D)	I+2		
192.	Stan	dard used for NMR is				
	(A).	methyl silane	(B)	triethyl silane		
	(C)	trimethyl silane	(b)	tetramethyl silane		
193.	In NMR spectroscopy the difference between the resonance position of a nucleus and that of					
	a sta	ndard reference compound is called as				
	(A)	spin spin interaction	(B)	proton magnetic resonance		
	(C)	spin spin coupling	S)	chemical shift		
194.	In N	MR spectroscopy the distance betwee	en the	e centres of the two adjacent peaks in a		
	1.1	iplet usually is constant and is called				
		Coupling constant	(B)	Spin rotation constant		
	(C)	Shift constant	(D)	Peak constant		
195.	Whic	h of the following equipment is require	ed for c	conducting radio immuno array?		
	(A)	pH meter	(B)	Centrifuge		
	(C)	Conductometer	(D)	Densitometer		

196.	In wh	ich of the following ways the capillary electrophoretic separations are performed?
	(A)	Iso electric focussing
	(B)	Fast atom bombardment
	(C)	Double focussing spectrometers
	(D)	Field ionization
197.	Cond	uctance is expressed as ——— units.
	(A)	A° ohms ⁻¹
	(C)	δ (D) MeV
198.	Ampe	erometric titrations are performed using — method.
	(A)	droping mercury electrode
	(B)	glass electrode
	(C)	polarographic
	(D)	specific ion electrode
199.	Stret	ching vibration in IR spectroscopy involves changes in the
	(A)	bond angle bond length
	(C)	bond rotation (D) bond bending
200.	Stror	nger bonds produce IR absorption at higher frequencies which of the following would
	be?	
	(A)	sp (B) sp ²
	(C)	$\mathrm{sp^3}$ (D) $\mathrm{sp^4}$

SPACE FOR ROUGH WORK

SPACE FOR ROUGH WORK

SPACE FOR ROUGH WORK



DJPC/19 40